

Vacuum-ultraviolet interconfigurational $4f^3 \rightarrow 4f\ 25d$ absorption and emission studies of the Nd^{3+} ion in KYF, YF, and YLF crystal hosts

Kollia Z., Sarantopoulou E., Cefalas A., Nicolaides C., Naumov A., Semashko V., Abdulsabirov R., Korableva S., Dubinskii M.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The laser-induced vacuum-ultraviolet fluorescence spectra of $\text{KY } 3\text{F}_{10}:\text{Nd}^{3+}$ (KYF:Nd) and $\text{YF}_3:\text{Nd}^{3+}$ (YF:Nd) single crystals pumped by a pulsed-discharge molecular F2 laser at 157 nm were obtained. A number of new fluorescence peaks were observed and were assigned to the $4f\ 25d \rightarrow 4f^3$ dipole-allowed transitions of the Nd^{3+} ion. The absorption spectra of $\text{LiYF}_4:\text{Nd}^{3+}$ (YLF:Nd), KYF:Nd, and YF:Nd crystal samples in the vacuumultraviolet spectral regions were also obtained. Finally, the splitting of the states of the $4f25d$ configuration of the Nd^{3+} ion, which is due to the crystal field, was observed in all the crystal samples. © 1995 Optical Society of America.

<http://dx.doi.org/10.1364/JOSAB.12.000782>
